

Take Control To The Third Power



PoultrySulfa™ – The *Only* FDA-Approved Triple-Sulfa Soluble Powder Combination

Fight coccidiosis and fowl cholera with the unique broad-spectrum “triple-sulfa” protection of PoultrySulfa and take control *to the third power*.

- Three active ingredients in one for broad-spectrum control
- Combination of sulfas gives increased effectiveness and low toxicity
- Clearly soluble

Indications

As an aid in the control of coccidiosis and acute fowl cholera in chickens and acute fowl cholera and coccidiosis in turkeys, when caused by pathogens susceptible to sulfamethazine, sulfamerazine, and sulfaquinoxaline.

Composition (each pack contains)

78 g sodium sulfamethazine activity
78 g sodium sulfamerazine activity
39 g sodium sulfaquinoxaline activity

PoultrySulfa™

Soluble Powder

PoultrySulfa Soluble Powder

Dosage And Administration

Disease	Treatment
Acute Fowl Cholera TURKEYS & CHICKENS	Provide medicated water (0.040% solution) for 2 to 3 days. If disease recurs, repeat treatment.
Coccidiosis TURKEYS	Provide medicated water (0.025% solution) for 2 days, then plain water for 3 days, then medicated water (0.025% solution) for 2 days, then plain water for 3 days, then medicated water (0.025% solution) for 2 days. Repeat if necessary. DO NOT CHANGE LITTER.
Coccidiosis CHICKENS	Provide medicated water (0.040% solution) for 2 to 3 days, then plain water for 3 days, then medicated water (0.025% solution) for 2 days. If bloody droppings appear, repeat at 0.025% level for 2 more days. DO NOT CHANGE LITTER.

Proportioner Mixing Instructions

Add the following amount of water to one pack of **PoultrySulfa** to make a stock solution for proportioner set to meter at the rate of one ounce per gallon (1:128).
Make fresh solutions daily.

Concentration in drinking water	Amount of water to add to one pack to make a stock solution for proportioner set at 1:128
0.040%	1 gallon (3.8 liters)
0.025%	1.6 gallons (6.1 liters)

Tank Mixing Instructions

Add one pack to a tank of adequate volume and then add the amount of water listed below to form the treatment solution. Make fresh solutions daily.

Concentration in drinking water	Amount of water per one pack
0.040%	128 gallons
0.025%	206 gallons

WARNING (HUMAN FOOD)

DO NOT TREAT CHICKENS OR TURKEYS WITHIN 14 DAYS OF SLAUGHTER FOR FOOD. Do not medicate chickens or turkeys producing eggs for human consumption. For oral veterinary use only. Keep all medications out of the reach of children. Dispose any waste and unused portions properly. SEE PACKAGE LABEL FOR COMPLETE USE DIRECTIONS AND WARNINGS.

CAUTION

If improvement is not noted in 72 hours, consult a poultry veterinarian or poultry diagnostic laboratory for diagnosis. During treatment use only medicated water unless otherwise directed. May cause toxic reactions unless drug is evenly mixed in water at dosages indicated and used according to label instructions. For control of disease outbreaks medication should be initiated as soon as diagnosis is determined. Treated animals must actually consume enough medicated water to provide a therapeutic dose. Do not mix or administer in galvanized containers.

For more information on putting PoultrySulfa to work for you, contact your nearest Alpharma representative.



Alpharma Inc.
One Executive Drive
Fort Lee, NJ 07024, USA
(800) 643-5791 www.alpharma.com

PoultrySulfa is a trademark of Alpharma Inc.
©2005 Alpharma Inc. All rights reserved. PD0512

PoultrySulfa™
Soluble Powder